Half-life for grafts survival one year (years) **C** 20 30 75 76 77 78 79 80 81 82 83 84 85 86 87 88 89 90 91 92 93 94 95 Half-life after 1 year The first water made in the first the first term term term term from the first One-year survival .**:**5-Figure 20 T 0.8 100 60 One year survival (%)


Figure 2

EFFECT OF TGF-B ON VARIOUS T CELL SUBSETS IN THE GENERATION OF SUPPRESSORS OF CYTOTOXIC T CELL ACTIVITY

Effector: Target Ratio CD8 CD45RA ---- Allo-CTL activity → + TGFβ CD8 ---+ Con CD8 25-507 Effector: Target ratio CD4 CD45RO ---- Allo-CTL activity 50- -- + TGFβ CD4 -o- + Con CD4 20 25-7 9 Effector: Target ratio CD4 CD45RA ---- Allo-CTL activity 50- -- + TGFβ CD4 --- + Con CD4 25sisyJ %

EFFECT OF CD4 CELLS PRIMED WITH TGF-β (CD4reg) ON ALLO-CTL ACTIVITY






100

Figure 4

grate that the treet much them then the sense of the state of the stat

CD4 REGULATORY T CELLS REQUIRE CELL CONTACT TO INHIBIT CTYTOXIC T LYMPHOCYTE ACTIVITY

With Effector Cells



Separated from Effector cells




FIGURE 5





Figure 6






Figure 7

